

Jassa marmorata

Tube dwelling amphipod

Threat scores

- i. Ecological impact
 - Can compete with native marine organisms for food and space
 - Studies have also shown that Amphipods may play important roles in determining the type and distribution of algal communities, particularly where predation pressure is low (Molnar 2008)
- 2. Invasive potential
 - Juveniles are responsible for short distance dispersal
 - Low potential for local dispersal of adults once inoculated into an area, they tend to remain a local invasive species at the point of inoculation (Molnar 2008)
- 3. Geographic extent
 - Locally pervasive
- 4. Management difficulty
 - Little info on control and management
 - Ballast treatment may be key to preventing long distance spread

Geography and Habitat

- 1. Origin: North Atlantic Ocean and the Mediterranean Sea
- 2. Introduced: Alaska to California
- Habitats
 - Marine, fouling communities

Invasion Pathways

- 1. Ballast Water and Sediments
- 2. Hull/Surface Fouling
- 3. Natural Spread

Non native locations

- 1. 54- Gulf of Alaska
- 2. 56- Puget Trough/Georgia Basin
- 3. 58- Northern California

Sources

- 1. Molnar, Jennifer, et al. 2008. "Assessing the global threat of invasive species to marine biodiversity." *Frontiers in Ecology and the Environment*. 6 (9), pp. 485-492.
- 2. http://conserveonline.org/workspaces/global.invasive.assessment
- 3. http://nas.er.usgs.gov/XIMAGESERVERX/2009/20090409151343.jpg

